



The M-CATs

SEMINAR 8

Table of Contents

CH 10

General
Chemistry
Acid Base Chemistry


CH 11

General
Chemistry
Redox Reactions




General Chemistry Chapter 10

Acids & Bases





Basic Definition

Acid:
(+)



Base:
(-)



Arrhenius Theory

Acid: produces H^+ (aq)



HCl, H₂SO₄, HClO₄, HF...

Base: produces OH^- (aq)



NaOH, Ca(OH)₂, KOH...





Brønsted-Lowry Theory

Acid: H^+ donor

Base: H^+ acceptor




10.1: Definitions




Lewis Theory

Acid: e⁻ acceptor


Base: e⁻ donor



10.1: Definitions



Autoionization of Water



$$\begin{aligned} K_w &= [H_3O^+][OH^-] \\ &= (10^{-7}M)(10^{-7}M) = 10^{-14} \text{ (25}^\circ\text{C)} \end{aligned}$$

$$pH = -\log[H^+] = \log(1/[H^+])$$

$$pOH = -\log[OH^-] = \log(1/[OH^-])$$

$$pH + pOH = 14$$

10.2: Properties



Question

(a). An unknown powder is dissolved in 2.0L of water at standard ambient temperature and pressure. The resulting solution contains 0.0035 mol of H^+ .

What is the pH of the solution?

- A. 3.4
- B. 2.8
- C. 2.2
- D. 6.7

(b)What is the pOH?

- A. 11.2
- B. 10.5
- C. 10.8
- D. 6.7



Ionization of Acids and Bases (aq)

Acid:



$$K_a = \frac{[A^-][H_3O^+]}{[HA]}$$

Base:



$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$pH + pOH = 14 \text{ @ SATP}$$

10.2: Properties



Strong Acids and Bases (aq)

$$K_a \text{ or } K_b \gg 1$$



Acid:

$$\text{IF } [HA] \gg 10^{-7} M: \quad pH = -\log[HA]$$

$$\text{IF } [HA] \approx 10^{-7} M: \quad pH = -\log([x + HA])$$

$$K_w = [H^+][OH^-] = [HA + x][x] = 10^{-14}$$

HCl, HBr, HI, HNO₃, H₂SO₄, HClO₄, HClO₃

Base:

$$\text{IF } [B] \gg 10^{-7} M: \quad pOH = -\log[B]$$

$$\text{IF } [B] \approx 10^{-7} M: \quad pH = -\log([x + B])$$

$$K_w = [H^+][OH^-] = [x][B + x] = 10^{-14}$$

NaOH, KOH, Ca(OH)₂, Ba(OH)₂...



Question

A student prepares a solution by adding 2.0 mL of 0.010 M HCl to 998 mL of distilled water at 25°C. Part

(a) What is the pH of the resulting solution?

- A. 3.72
- B. 4.70
- C. 8.68
- D. 6.70

The student performs a 1:100 dilution using 10 mL of the original solution under the same conditions.

(b) What is the pH of the resulting solution?

- A. 5.45
- B. 6.62
- C. 6.80
- D. 6.95




Weak Acids and Bases (aq)

Acid:




$$K_a = \frac{[A^-][H_3O^+]}{[HA]} < 1$$



Base:



$$K_b = \frac{[OH^-][HB^+]}{[B]} < 1$$



$$K_a * K_b = K_w$$

$$\therefore pK_a + pK_b = 14$$

10.2: Properties



Question

A student prepares a 0.50 M solution of HF in water. The acid dissociation constant for HF at 25°C is:


$$K_a = 6.8 \times 10^{-4}$$

What is the solution's pH?

- A. 1.0
- B. 1.7
- C. 2.3
- D. 3.2



Neutralization




A strength	B strength	Product(s)	pH @ eq	E.g.
Strong	Strong	Salt+H ₂ O	7	HCl(aq)+NaOH(aq)-> NaCl(aq)+H ₂ O(l)
Strong	Weak	Salt	<7	HBr (aq)+NH ₃ (aq)-> NH ₄ Br(aq)
Weak	Strong	Salt+H ₂ O	>7	HF(aq)+KOH(aq)->KF(aq)+H ₂ O(l)
Weak	Weak	Salt	Idk	HCOOH(aq)+C ₅ H ₅ N(aq)->C ₅ H ₅ NH ⁺ HCOO ⁻ (aq)

10.2: Properties




Acid-Base Titrations



Curves

A strength	B strength	pH @ eq
Strong	Strong	7
Strong	Weak	<7
Weak	Strong	>7
Weak	Weak	Idk



Henderson-Hasselbalch Equation

$$pH = pK_a + \log \left[\frac{A^-}{HA} \right]$$

$$pOH = pK_b + \log \left[\frac{BH^{+}}{B} \right]$$



Question

What is the pH of a solution made from 1 L of 0.05 *M* acetic acid (CH_3COOH , $K_a = 1.8 \times 10^{-5}$) mixed with 500 mL of 1 *M* acetate (CH_3COO^-)?

- a. 3.9
- b. 4.5
- c. 5.2
- d. 5.7



Question

A diprotic acid H_2A has:

$$pK_{a1} = 3.0, pK_{a2} = 7.0$$


During titration with a strong base, the solution reaches the second equivalence point. Assume 25°C. What is the approximate pH at the second equivalence point?

- A. 3.0
- B. 5.0
- C. 9.0
- D. 11.0



General Chemistry Chapter 11

Redox Reactions



Redox Basics




*11.1 Oxidation-
Reduction Rxns*

Good OAs: O_2 , H_2O_2 , X_2 , H_2SO_4 , HNO_3 , $NaClO$, $KMnO_4$, CrO_3 , $NaCr_2O_7$, PCC, NAD^+ , FADH

Good RAs: CO , C , B_2H_6 , Sn^{2+} , M^0 s, hydrazine, Zn/Hg , Lindlar's cat., $NABH_4$, $LiAlH_4$, $NADH$, $FADH_2$

Question

#X. The following molecule is often used in redox reactions. What is the most likely reason for this?



- A. It is a reducing agent
- B. It is an oxidizing agent
- C. It can serve both as an electron acceptor or donor, catalyzing a redox reaction
- D. It is a precursor to an oxidizing agent



Redox Half Reactions

1. Determine oxidation numbers
2. Split equations into half rxns
3. Balance all except H and O
4. Balance O and H per environment
5. Balance e^-

*11.1 Oxidation-
Reduction Rxns*



Net Ionic Equations


For Redox:

1. Single Displacement
2. Combination
3. Decomposition
4. Combustion
5. Disproportionation

*11.2: Net Ionic
Equations*



Redox Titrations



11.3: Titrations



Question

A 25.00 mL sample containing Fe^{2+} is titrated with 0.02000 M $KMnO_4$ in acidic solution to produce Fe^{3+} and Mn^{2+} . It requires 18.60 mL of the $KMnO_4$ solution to reach the endpoint.

The balanced reaction is:

What is the molarity of iron (II) in the sample in M?

- a. 0.06700
- b. 0.07440
- c. 0.00850
- d. 0.03260

